STOP! READ ENTIRE LABEL BEFORE EACH USE.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SHAKE WELL BEFORE USING.

Use Restrictions • Except when applying to dogs, do not allow adults, children or pets to enter the area until sprays have dried. • Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

DOG, DOG BEDDING AND PREMISES: USE ONLY ON DOGS: To control fleas, ticks and lice: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and under body. While spraying, fluff the hair so the spray will penetrate to the skin. Make sure spray west thoroughly, but do not saturate animal. Do not spray into eyes or face. Do not contact with genitalia. Use protective gloves or mitts when applying product. Repeat application every two weeks if necessary. DO NOT USE ON CATS DR RITTENS. Do not use on dogs under twelve (12) weeks old. Consult a veterinarian before using this product on medicated, debilitated, aged, pregnant or nursing animals. Sensitivities may occur after using ANY pesticide products for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amount of water. If signs continue, consult a veterinarian immediately.

To control fleas and ticks on dog bedding, premises and upholstered furniture: Fleas and ticks (adults and larvae):
Thoroughly clean or launder dog bedding before treatment. Spray bedding, sleeping quarters, floor coverings and floors, paying special attention to cracks and crevices. Upholstered furniture may also be treated to control fleas and ticks in these areas. A test application should be made to upholstery in an inconspicuous place before use. Spray upholstery, paying particular attention to tufts and seams. Allow to dry before reuse.

INDOOR APPLICATION: For spot application only. Do not use as a space spray indoors. In the home, all food-processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed. Do not use in commercial food/feed-handling establishments, restaurants or other establishments where food/feed is commercially prepared, processed or served.

To control fleas (adults and larvae) and ticks, including ticks that may carry Lyme Disease (adults and larvae):
Thoroughly spray infested areas, dog beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as infestation occurs, but not more than once every two weeks. Allow to dry thoroughly before allowing pets to contact treated areas.

STORAGE AND DISPOSAL

Pesticide Storage: Store in a cool, dry area, inaccessible to children and away from heat or open flame.

Pesticide Disposal and Container Handling: Do Not Puncture or Incinerate! If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.



DO NOT USE ON CATS. **May be toxic or potentially fatal if applied to or ingested by cats.** Accidental application to cats and/or grooming a recently treated dog may result in tremors and/or uncoordinated muscle movements. If this occurs, immediate veterinary care should be provided.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice.

Have the product container with you when calling a Poison Control Center or doctor, or going for treatment. You may also contact 1-800-917-5431 for emergency medical treatment information.

Physical or Chemical Hazards • Contents under pressure. Do not use or store near heat or open flame. Exposure to temperatures above 130°F may cause bursting.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Questions or Comments? Call 1-800-917-5431 or visit our website at www.hotshot.com

[†]**GUARANTEED:** If you are not satisfied with this product for any reason, send your written request for a refund to the address below, together with the dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

FPO UPC replace and verify readability 0 71121 02117 5

Distributed *by* Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, M0 63114-0642
EPA Reg. No. 305-55-8845
EPA Est. Nos. 9688-M0-1①, 3657-WI-2®U, 58996-M0-1@EPA Est. Nos. 9688-M0-1①, 3657-WI-2®U, 58996-M0-1@EPA Est. Nos. 90202 United Industries Corporation.
All Rights Reserved.





FLEA, TICK & LICE KILLER with Odor Neutralizer

Provides Long-Lasting Protection

One Treatment Lasts 2 Weeks

FOR TREATMENT OF CARPETING & UPHOLSTERED FURNITURE
• Can Be Used On Dogs & Dog Bedding

ACTIVE INGREDIENT:
Permethrin. 0.5%
OTHER INGREDIENTS 99.5%

Net Wt 14 oz (396 a)

KEEP OUT OF REACH OF CHILDREN DO NOT USE CAUTION See back to ON CATS.

*For terms of guarantee, see back.